Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

FOREIGN AND INSULAR STATISTICAL REPORTS OF COUNTRIES AND CITIES—UNTABULATED.

AUSTRIA—*Brünn.*—Month of May, 1906. Estimated population, 95,342. Total number of deaths, 230, including diphtheria 1, whooping cough 1, and 60 from tuberculosis.

BRAZIL—*Pernambuco.*—Two weeks ended May 31, 1906. Estimated population, 200,000. Total number of deaths, 363, including smallpox 5, and 60 from tuberculosis.

CANADA—*Ontario—Hamilton.*—Month of June, 1906. Estimated population, 59,567. Total number of deaths not reported. Two deaths from diphtheria, 1 from whooping cough, and 4 from tuberculosis reported.

Quebec—Sherbrooke.—Month of June, 1906. Estimated population, 13,950. Total number of deaths, 18, including 1 from enteric fever.

FRANCE—*Roubaix.*—Month of June, 1906. Estimated population, 124,660. Total number of deaths, 148, including measles 3 and 1 from whooping cough.

GIBRALTAR.—Three weeks ended July 1, 1906. Estimated population, 27,460. Total number of deaths, 38, including 2 from smallpox.

GREAT BRITAIN—*England and Wales.*—The deaths registered in 76 great towns in England and Wales during the week ended June 30, 1906, correspond to an annual rate of 12.4 per 1,000 of population, which is estimated at 15,818,360.

Bradford.—Two weeks ended June 30, 1906. Estimated population, 288,544. Total number of deaths, 163, including measles 10, whooping cough 1, and 17 from phthisis pulmonalis.

London.—One thousand and sixty deaths were registered during the week, including measles 46, scarlet fever 5, diphtheria 9, enteric fever 9, whooping cough 14, and 22 from diarrhea. The deaths from all causes correspond to an annual rate of 11.7 per 1,000. In Greater London 1,495 deaths were registered. In the "outer ring" the deaths included 5 from diphtheria, 8 from measles, 2 from scarlet fever, and 2 from whooping cough.

Ireland.—The average annual death rate represented by the deaths registered during the week ended June 30, 1906, in the 21 principal town districts of Ireland was 15.4 per 1,000 of the population, which is estimated at 1,093,959. The lowest rate was recorded in Drogheda, viz, 4.1, and the highest in Armagh, viz, 14.5 per 1,000. In Dublin and suburbs 125 deaths were registered, including enteric fever 2, whooping cough 1, and 30 from tuberculosis.

Scotland.—The deaths registered in 8 principal towns during the week ended June 30, 1906, correspond to an annual rate of 14.5 per 1,000 of the population, which is estimated at 1,787,788. The lowest rate of mortality was recorded in Aberdeen, viz, 11.9, and the highest

in Perth, viz, 21.1 per 1,000. The aggregate number of deaths registered from all causes was 497, including measles 11, diphtheria 6, enteric fever 10, scarlet fever 3, and 9 from whooping cough.

HAWAII—*Honolulu.*—Month of June, 1906. Estimated population, 39,306. Total number of deaths, 76, including diphtheria 1, plague 3, and 12 from tuberculosis.

JAPAN—*Nagasaki.*—Ten days ended June 20, 1906. Estimated population, 163,324. Total number of deaths not reported. One death from diphtheria and 1 from enteric fever reported.

SPAIN—*Barcelona.*—Ten days ended June 20, 1906. Estimated population, 600,000. Total number of deaths, 303, including diphtheria 1, enteric fever 4, measles 8, scarlet fever 1, smallpox 5, and 31 from tuberculosis.

Seville.—Month of May, 1906. Estimated population, 148,315. Total number of deaths, 464, including diptheria 4, enteric fever 3, whooping cough 2, smallpox 53, and 74 from tuberculosis.

WEST INDIES.—*Curaçao.*—Three weeks ended July 6, 1906. Estimated population, 31,600. Total lnumber of deaths, 15, including 1 from typhus fever.

Cholera, yellow fever, plague, and smallpox, from June 29 to July 27, 1906.

[Reports received by the Surgeon-General, Public Health and Marine-Hospital Service, from American consuls through the Department of State, and from other sources.]

[For reports received from December 29, 1905, to June 29, 1906, see PUBLIC HEALTH REPORTS for June 29, 1906.]

[NOTE.—In accordance with custom, the tables of epidemic diseases are terminated semiannually and new tables begun.]

CHOLERA.

Place.	Date.	Cases.	Deaths.	Remarks.
India:				
Bombay	May 30–June 19	.....	57	
Calcutta	May 20–June 9	.....	104	
Madras	June 2–15	.....	5	
Rangoon	May 27–June 2	.....	2	
Philippine Islands:				
Manila	May 13–June 9	6	6	
Provinces	Apr. 28–June 9	45	30	1 on schr. Florence S.
Siam:				
Bangkok	Feb. 3–Apr. 14	224	198	
Straits Settlements:				
Parak	May 9–12	64	34	
Singapore	Apr. 25–May 8	17	22	
Wellesley Province	Apr. 10–May 12	107	81	

YELLOW FEVER.

Costa Rica:				
Limon	June 20–25	1	.....	
Honduras:				
Ceiba	July 21–July 23	.....	.....	Epidemic.
Pimienta	June 12–July 5	14	.....	
Mexico:				
Veracruz: (Veracruz)	July 23	1	.....	Imported from Yucatan on ss. Sinaloa.
Yucatan (Merida)	June 10–July 7	34	22	
Peru:				
Callao	May 27–June 2	.....	1	On Br. ss. Chile from Ancon.